



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Tables of Pure Reciprocants to the Weight 8.

BY PROF. CAYLEY.

In the tabulation of Pure Reciprocants it is convenient to write $a = 1$; we thus have for all the reciprocants of a given weight a single column of literal terms which (as in the Seminvariant Tables) I arrange in alphabetical order AO , and the several reciprocants have then each of them its own column of numerical coefficients: the form of the table is thus similar to that of the seminvariant table, the only difference being that for reciprocants the final terms are not in general power-enders: as in the seminvariant table, the columns of the table are arranged *inter se* with their final terms in AO . As remarked in my paper, "Corrected Seminvariant Tables for the Weights 11 and 12," A. M. J., t. XIV (1892), pp. 195–200, it is not in every case the top term of a column which should be regarded as the initial term; but to the extent 8 to which the reciprocant tables are here carried this remark has no application.

I recall that the notation is the modified one employed by Halphen, and by Sylvester in his 12th and subsequent lectures, viz. a, b, c, d, \dots denote $\frac{1}{2} \frac{d^2 y}{dx^2}, \frac{1}{6} \frac{d^3 y}{dx^3}, \frac{1}{24} \frac{d^4 y}{dx^4}, \frac{1}{120} \frac{d^5 y}{dx^5}, \dots$ respectively. As already noticed, a is put $= 1$, but it is to be in the several terms restored in the proper powers so as to obtain for the reciprocant a homogeneous expression of a degree equal to the original degree of the final term; thus $d - 3bc + 2b^3$ is to be read as standing for $a^2 d - 3abc + 2b^3$.

The ultimate verification of the expression for a pure reciprocant consists (as is known) in its annihilation by the operator

$V = 2a^2 \partial_b + 5ab \partial_c + (6ac + 3b^3) \partial_a + (7ad + 7bc) \partial_e + (8ae + 8bd + 4c^3) \partial_f + \text{etc.},$
or say

$V = 2\partial_b + 5b\partial_c + (6c + 3b^3)\partial_a + (7d + 7bc)\partial_e + (8e + 8bd + 4c^3)\partial_f + \text{etc.};$

thus for the reciprocant $50e - 175bd + 28c^2 + 105b^2c$, the result obtained is

$2(-175d + 210bc) + 5b(56c + 105b^2) + (6c + 3b^2)(-175b) + (7d + 7bc)(50)$,
or collecting, this is

$$\begin{array}{l|l} d & -350 \qquad \qquad \qquad +350 \\ bc & +420 + 280 - 1050 + 350 \\ b^3 & \qquad \qquad +525 - 525 \end{array} \left| \begin{array}{l} \pm 350 \\ \pm 1050 \\ \pm 525 \end{array} \right.$$

$= 0$, as it should be.

The tables are

c	+4
b^2	-5

+4
-5

d	+1
bc	-3
b^3	+2

± 3

e	+ 50	
bd	-175	
c^2	+ 28	+16
b^2c	+105	-40
b^4		+25

+183 +41
-175. -40

f	+10	
be	-40	
cd	-12	+ 4
b^2d	+65	- 5
bc^2	+16	-12
b^3c	-39	+23
b^5		-10

$\pm 91 \pm 27$

g	+ 14			
bf	- 63			
ce	-1350	+ 800		
d^2	+1470	- 875	+ 125	
b^2e	+1782	-1000		
bcd	-4158	+2450	- 750	
c^3	+2130	-1344	+ 256	+ 64
b^3d			+ 500	
b^2c^2		+ 35	+ 165	- 240
b^4c			- 300	+ 300
b^6				- 125

+5576 +3250 \pm 1018 + 364
-5508 -3254 - 365

h	+ 7			
bg	- 35			
cf	- 539	+ 560		
de	+ 605	- 650	+ 50	
b^2f	+ 735	- 700		
bce	+ 306	- 290	- 150	
bd^2	-2135	+2275	- 175	
c^2d	+1001	-1036	+ 28	+ 16
b^3e	-1485	+1500	+ 100	
b^2cd	+3465	-3710	+ 630	- 40
bc^3	-1295	+1988	- 84	- 48
b^4d			- 350	+ 25
b^3c^2		+ 63	- 259	+152
b^5c			+ 210	-155
b^7				+ 50

+6119 \pm 6386 \pm 1018 \pm 243
-5489

i	+	420							
bh	—	2310							
cg	—	32704	+	1176					
df	+	57750	—	8085	+	20433			
e^2	—	20460	+	7040	—	21542	+	625	
b^2g	+	45500	—	1470					
bcf	—	28392	+	18963	—	61299			
bde	—	90900	—	16940	+	69062	—	4375	
c^2e	+	103740	—	27160	+	80248	—	49700	
cd^2	—	38320	+	26460	—	85554	+	55125	
b^3f	—	69615	—	9555	+	40866			
b^3ce	+	83538	+	28098	—	106218	+	128625	
b^3d^2	+	92820	+	12740	—	54782	—	61250	
bc^2d	—	102102	—	52822	+	191590	—	156800	
c^4			+	21560	—	73304	+	84868	
b^4e					—	378	—	78750	
b^3cd					+	1176	+	183750	
b^2c^3							—	102165	
b^5d									
b^4c^2							+	175	
b^6c							+	1500	
b^8								+	625

$$\begin{aligned}
&+383768 + 116037 + 403375 + 452993 + 29130 + 8774 + 3281 \\
&-384803 - 116032 - 403077 - 453040 - 29126 - 8750 - 3280
\end{aligned}$$

I remark that in the last of these tables the first column, say $i \propto bc^2d$, which ends in bc^2d , is a more simple form than Sylvester's P_8 , $= i \propto c^4$, (A. M. J., t. IX, p. 35) which ends in c^4 ; P_8 is in fact a linear combination, first col. + 6 second col. of the first and second columns of the table: the second column, say $cg \propto c^4$ is Sylvester's (a^2cg), t. IX, p. 124.